

UDC 517.946

A. Gromyk*, Ph. D. of Tech. Sci,**I. Konet****, Doctor of Phys. and Mathem. Sciences, Professor,**T. Pylypiuk****, Ph. D of Physics and Mathem.*State Agrarian and Engineering University in Podilya,
Kamianets-Podilsky,**Kamianets-Podilsky Ivan Ohienko National University,
Kamianets-Podilsky

HYPERBOLIC BOUNDARY VALUE PROBLEM FOR SEMIBOUNDED PIECEWISE-HOMOGENEOUS HOLLOW CYLINDER

By means of the method of integral and hybrid integral transforms, in combination with the method of main solutions (influence matrices and Green matrices) the integral image of exact analytical solution of hyperbolic boundary value problem of mathematical physics for semibounded piecewise-homogeneous hollow cylinder is obtained for the first time.

Keywords: *hyperbolic equation, initial and boundary conditions, conjugate conditions, integral transforms, the main solutions.*

Introduction. The theory of hyperbolic boundary value problems for partial differential equations and including equations of mathematical physics, is an important section of the modern theory of differential equations which is intensively developing in the present time. The achievements of this theory apply to research of different mathematical models of various processes and phenomena of mechanics, physics, technology and new technologies.

Significant results from the theory of Cauchy and initial-boundary value problems (mixed problems) for hyperbolic equations were obtained in the known works of J. Hadamard [1], L. Gording [2], Yu. Mitropolsky, G. Khoma, M. Hromyak [3], A. Samoilenko, B. Tkach [4], M. Smirnov [5], V. Chernyatyn [6] and others domestic and foreign mathematics.

It is well known that the complexity of a boundary-value problem significantly depends on the coefficients of equations (different types of degeneracy and features) and the geometry of domain (limited, unlimited, smoothness of the boundary, the presence of corner points, etc.) in which the problem is considered. The dependence of the properties of solutions of boundary value problems for linear, quasi-linear, and certain classes of nonlinear equations (elliptic, parabolic, hyperbolic) in homogeneous domains (homogeneous environments) on the above-mentioned properties of the coefficients of equations and geometry of domain are studied in detail, and functional spaces of correctness of problems in the sense of Hadamard are constructed.

However, many important applied problems of thermomechanics, thermal physics, diffusion, theory of elasticity, theory of electrical circuits, theory of vibrations of mechanical systems lead to boundary value problems and mixed problems not only in homogeneous domains when the coefficients of the equations are continuous, but also in inhomogeneous and piecewise homogeneous domains when the coefficients of the equations are piecewise continuous or piecewise constant [7, 8].

The method of hybrid integral transforms generated by hybrid differential operators when in each component of connectivity of piecewise homogeneous domain are treated different differential operators or differential operators look the same, but with different sets of coefficients is an effective method of constructing exact solutions for a fairly broad class of linear boundary value problems and mixed problems in piecewise homogeneous domains [9–12].

This article is a logical extension of [13]. By means of the method of hybrid integral transforms the exact solution of hyperbolic boundary value problem of mathematical physics for semibounded piecewise homogeneous hollow cylinder is obtained in this article.

Formulation of the problem. Let's consider the problem of structure of 2π -periodic for angular variable φ solution of partial differential equations of hyperbolic type of 2nd order [14]

$$\frac{\partial^2 u_j}{\partial t^2} - \left[a_{rj}^2 \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} \right) + \frac{a_{\varphi j}^2}{r^2} \frac{\partial^2}{\partial \varphi^2} + a_{zj}^2 \frac{\partial^2}{\partial z^2} \right] u_j + \chi_j^2 u_j = f_j(t, r, \varphi, z); \quad r \in I_j; \quad j = \overline{1, n+1} \quad (1)$$

which is bounded in the set

$$D = \{(t, r, \varphi, z) : t > 0; r \in I_n^+ = \bigcup_{j=1}^{n+1} I_j \equiv \bigcup_{j=1}^{n+1} (R_{j-1}; R_j), R_0 > 0, R_{n+1} \equiv R < +\infty; \varphi \in [0; 2\pi); z \in (0; +\infty)\}$$

with initial conditions

$$u_j|_{t=0} = g_j^1(r, \varphi, z); \quad \frac{\partial u_j}{\partial t}|_{t=0} = g_j^2(r, \varphi, z); \quad r \in I_j; \quad j = \overline{1, n+1} \quad (2)$$

boundary conditions

$$\left(\alpha_{11}^0 \frac{\partial}{\partial r} + \beta_{11}^0 \right) u_1 \Big|_{r=R_0} = g_0(t, \varphi, z);$$

$$\left(\alpha_{22}^{n+1} \frac{\partial}{\partial r} + \beta_{22}^{n+1} \right) u_{n+1} \Big|_{r=R} = g(t, \varphi, z); \quad (3)$$

$$\left(-\frac{\partial}{\partial z} + h \right) u_j \Big|_{z=0} = g_j(t, r, \varphi); \quad \frac{\partial^s u_j}{\partial z^s} \Big|_{z=+\infty} = 0; \quad s = 0, 1; \quad j = \overline{1, n+1} \quad (4)$$

and conjugate conditions

$$\left[\left(\alpha_{j1}^k \frac{\partial}{\partial r} + \beta_{j1}^k \right) u_k - \left(\alpha_{j2}^k \frac{\partial}{\partial r} + \beta_{j2}^k \right) u_{k+1} \right]_{r=R_k} = 0; \quad j = 1, 2; \quad k = \overline{1, n}, \quad (5)$$

here $a_{rj}, a_{\varphi j}, a_{zj}, \chi_j, h, \alpha_{js}^k, \beta_{js}^k$ — some constants;

$$\alpha_{11}^0 \leq 0; \beta_{11}^0 \geq 0; \left| \alpha_{11}^0 \right| + \beta_{11}^0 \neq 0; \quad \alpha_{22}^{n+1} \geq 0; \beta_{22}^{n+1} \geq 0; \alpha_{22}^{n+1} + \beta_{22}^{n+1} \neq 0;$$

$$c_{jk} = \alpha_{2j}^k \beta_{1j}^k - \alpha_{1j}^k \beta_{2j}^k \neq 0; \quad c_{1k} \cdot c_{2k} > 0;$$

$$f(t, r, \varphi, z) = \{ f_1(t, r, \varphi, z), f_2(t, r, \varphi, z), \dots, f_{n+1}(t, r, \varphi, z) \};$$

$$g^1(r, \varphi, z) = \{ g_1^1(r, \varphi, z), g_2^1(r, \varphi, z), \dots, g_{n+1}^1(r, \varphi, z) \};$$

$$g^2(r, \varphi, z) = \{ g_1^2(r, \varphi, z), g_2^2(r, \varphi, z), \dots, g_{n+1}^2(r, \varphi, z) \};$$

$$g(t, r, \varphi) = \{ g_1(t, r, \varphi), g_2(t, r, \varphi), \dots, g_{n+1}(t, r, \varphi) \};$$

$$g_0(t, \varphi, z), \quad g(t, \varphi, z)$$

are known bounded continuous functions;

$$u(t, r, \varphi, z) = \{ u_1(t, r, \varphi, z), u_2(t, r, \varphi, z), \dots, u_{n+1}(t, r, \varphi, z) \}$$

is limited and twice-continuously differentiable desired function.

The main part. Let's assume that the solution of the problem (1)–(5) exists and defined and the unknown functions satisfy the condition of applicability of direct and inverse integral and hybrid integral transformations [15–17].

Let's apply the integral Fourier transform on Cartesian semiaxis $(0; +\infty)$ relative to variable z to the initial-boundary problem (1)–(5) [15]:

$$F_+ [f(z)] = \int_0^{+\infty} f(z) K(z, \sigma) dz \equiv \tilde{f}(\sigma), \quad (6)$$

$$F_+^{-1} [\tilde{f}(\sigma)] = \int_0^{+\infty} \tilde{f}(\sigma) K(z, \sigma) d\sigma = f(z), \quad (7)$$

$$F_+ \left[\frac{d^2 f}{dz^2} \right] = -\sigma^2 \tilde{f}(\sigma) + K(0, \sigma) \left(-\frac{df}{dz} + hf \right) \Big|_{z=0}, \quad (8)$$

here conversion kernel is $K(z, \sigma) = \sqrt{\frac{2}{\pi}} \frac{\cos(\sigma z) + h \sin(\sigma z)}{\sqrt{\sigma^2 + h^2}}$.

The integral operator F_+ due to the formula (6) as a result of identity (8) three-dimensional initial boundary value problem of conjugation (1)–(5) puts in accordance the task of constructing solution which is limited in the set $D' = \{ (t, r, \varphi); t > 0; r \in I_n^+; \varphi \in [0; 2\pi) \}$ and is 2π -periodical of angular variable φ of differential equations

$$\frac{\partial^2 \tilde{u}_j}{\partial t^2} - \left[a_{rj}^2 \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} \right) + \frac{a_{\varphi j}^2}{r^2} \frac{\partial^2}{\partial \varphi^2} \right] \tilde{u}_j + (a_{zj}^2 \sigma^2 + \chi_j^2) \tilde{u}_j = \quad (9)$$

$$= \tilde{F}_j(t, r, \varphi, \sigma); r \in I_j; j = \overline{1, n+1},$$

with initial conditions

$$\tilde{u}_j \Big|_{t=0} = \tilde{g}_j^1(r, \varphi, \sigma); \quad \frac{\partial \tilde{u}_j}{\partial t} \Big|_{t=0} = \tilde{g}_j^2(r, \varphi, \sigma); \quad r \in I_j; j = \overline{1, n+1}; \quad (10)$$

boundary conditions

$$\left(\alpha_{11}^0 \frac{\partial}{\partial r} + \beta_{11}^0 \right) \tilde{u}_1 \Big|_{r=R_0} = \tilde{g}_0(t, \varphi, \sigma); \quad \left(\alpha_{22}^{n+1} \frac{\partial}{\partial r} + \beta_{22}^{n+1} \right) \tilde{u}_{n+1} \Big|_{r=R} = \tilde{g}(t, \varphi, \sigma); \quad (11)$$

and conjugate conditions

$$\left[\left(\alpha_{j1}^k \frac{\partial}{\partial r} + \beta_{j1}^k \right) \tilde{u}_k - \left(\alpha_{j2}^k \frac{\partial}{\partial r} + \beta_{j2}^k \right) \tilde{u}_{k+1} \right] \Big|_{r=R_k} = 0; \quad j = 1, 2; \quad k = \overline{1, n}, \quad (12)$$

here $\tilde{F}_j(t, r, \varphi, \sigma) = \tilde{f}_j(t, r, \varphi, \sigma) + a_{zj}^2 K(0, \sigma) g_j(t, r, \varphi); \quad j = \overline{1, n+1}.$

Let's apply finite integral Fourier transform relative to the variable φ to the problem (9)–(12) [16]:

$$F_m [g(\varphi)] = \int_0^{2\pi} g(\varphi) \exp(-im\varphi) d\varphi \equiv g_m, \quad i = \sqrt{-1}; \quad (13)$$

$$F_m^{-1}[g_m] = \frac{\text{Re}}{2\pi} \sum_{m=0}^{\infty} \varepsilon_m g_m \exp(im\varphi) \equiv g(\varphi), \quad (14)$$

$$F_m \left[\frac{d^2 g}{d\varphi^2} \right] = -m^2 F_m [g(\varphi)] \equiv -m^2 g_m, \quad (15)$$

here $\text{Re}(\dots)$ – the real part of the expression (\dots) relative to the variable φ ; $\varepsilon_0 = 1, \quad \varepsilon_k = 2; \quad k = 1, 2, 3, \dots$

The integral operator F_m due to the formula (13) as a result of identity (15) two-dimensional initial boundary value problem of conjugation (9)–(12) puts in accordance the task of constructing solution which is limited in the set $D'' = \{(t, r); t > 0; r \in I_n^+\}$ of one-dimensional hyperbolic differential equations of 2nd order

$$\frac{\partial^2 \tilde{u}_{jm}}{\partial t^2} - a_{rj}^2 \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} - \frac{v_{jm}^2}{r^2} \right) \tilde{u}_{jm} + (a_{zj}^2 \sigma^2 + \chi_j^2) \tilde{u}_{jm} = \quad (16)$$

$$= \tilde{F}_{jm}(t, r, \sigma); \quad r \in I_j; \quad j = \overline{1, n+1}; \quad v_{jm} = a_{\varphi j} m / a_{rj}$$

with initial conditions

$$\tilde{u}_{jm}|_{t=0} = \tilde{g}_{jm}^1(r, \sigma); \quad \left. \frac{\partial \tilde{u}_{jm}}{\partial t} \right|_{t=0} = \tilde{g}_{jm}^2(r, \sigma); \quad r \in I_j; \quad j = \overline{1, n+1}, \quad (17)$$

boundary conditions

$$\left(\alpha_{11}^0 \frac{\partial}{\partial r} + \beta_{11}^0 \right) \tilde{u}_{1m} \Big|_{r=R_0} = \tilde{g}_{0m}(t, \sigma); \quad \left(\alpha_{22}^{n+1} \frac{\partial}{\partial r} + \beta_{22}^{n+1} \right) \tilde{u}_{n+1,m} \Big|_{r=R} = \tilde{g}_m(t, \sigma); \quad (18)$$

and conjugate conditions

$$\left[\left(\alpha_{j1}^k \frac{\partial}{\partial r} + \beta_{j1}^k \right) \tilde{u}_{km} - \left(\alpha_{j2}^k \frac{\partial}{\partial r} + \beta_{j2}^k \right) \tilde{u}_{k+1,m} \right] \Big|_{r=R_k} = 0; \quad j = 1, 2; \quad k = \overline{1, n}. \quad (19)$$

Let's apply finite hybrid integral Hankel transform of 2nd kind relative to the variable r in piecewise homogeneous segment I_n^+ of n conjugation points to the problem (16)–(19) [17]:

$$H_{sn} [f(r)] = \int_{R_0}^R f(r) V(r, \lambda_s) \sigma(r) r dr \equiv \tilde{f}(\lambda_s), \quad (20)$$

$$H_{sn}^{-1} [\tilde{f}(\lambda_s)] = \sum_{s=1}^{\infty} \tilde{f}(\lambda_s) \frac{V(r, \lambda_s)}{\|V(r, \lambda_s)\|^2} \equiv f(r), \quad (21)$$

$$\begin{aligned} H_{sn} [B_{(m)}[f(r)]] = & -\lambda_s^2 \tilde{f}(\lambda_s) - \sum_{k=1}^{n+1} \gamma_k^2 \int_{R_{k-1}}^{R_k} f(r) V_k(r, \lambda_s) \sigma_k r dr - \\ & - a_1^2 R_0 \sigma_1 \left(\alpha_{11}^0 \right)^{-1} V_1(R_0, \lambda_s) \left(\alpha_{11}^0 \frac{df}{dr} + \beta_{11}^0 f \right) \Big|_{r=R_0} + \\ & + a_{n+1}^2 R \sigma_{n+1} \left(\alpha_{22}^{n+1} \right)^{-1} \left(\alpha_{22}^{n+1} \frac{df}{dr} + \beta_{22}^{n+1} f \right) \Big|_{r=R}. \end{aligned} \quad (22)$$

Spectral function $V(r, \lambda_s)$, weight function $\sigma(r)$ and hybrid Bessel differential operator $B_{(m)} = \sum_{k=1}^{n+1} a_{rk}^2 \theta(r - R_{k-1}) \theta(R_k - r) B_{V_{km}}$, written in [17], take part in formulas (20)–(22).

Here $B_{V_{km}} = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} - \frac{V_{km}^2}{r^2}$ is classical Bessel differential operator, $\theta(x)$ is the Heaviside step function.

Let's write the differential equations (16) and the initial conditions (17) in matrix form

$$\begin{bmatrix} \left(\frac{\partial^2}{\partial t^2} - a_{r1}^2 B_{V_{1m}} + q_1^2(\sigma)\right) \tilde{u}_{1m} \\ \left(\frac{\partial^2}{\partial t^2} - a_{r2}^2 B_{V_{2m}} + q_2^2(\sigma)\right) \tilde{u}_{2m} \\ \dots\dots\dots \\ \left(\frac{\partial^2}{\partial t^2} - a_{r,n+1}^2 B_{V_{n+1,m}} + q_{n+1}^2(\sigma)\right) \tilde{u}_{n+1,m} \end{bmatrix} = \begin{bmatrix} \tilde{F}_{1m}(t, r, \sigma) \\ \tilde{F}_{2m}(t, r, \sigma) \\ \dots\dots\dots \\ \tilde{F}_{n+1,m}(t, r, \sigma) \end{bmatrix}, \quad (23)$$

$$\begin{bmatrix} \tilde{u}_{1m}(t, r, \sigma) \\ \tilde{u}_{2m}(t, r, \sigma) \\ \dots\dots\dots \\ \tilde{u}_{n+1,m}(t, r, \sigma) \end{bmatrix}_{t=0} = \begin{bmatrix} \tilde{g}_{1m}^1(r, \sigma) \\ \tilde{g}_{2m}^1(r, \sigma) \\ \dots\dots\dots \\ \tilde{g}_{n+1,m}^1(r, \sigma) \end{bmatrix};$$

$$\frac{\partial}{\partial t} \begin{bmatrix} \tilde{u}_{1m}(t, r, \sigma) \\ \tilde{u}_{2m}(t, r, \sigma) \\ \dots\dots\dots \\ \tilde{u}_{n+1,m}(t, r, \sigma) \end{bmatrix}_{t=0} = \begin{bmatrix} \tilde{g}_{1m}^2(r, \sigma) \\ \tilde{g}_{2m}^2(r, \sigma) \\ \dots\dots\dots \\ \tilde{g}_{n+1,m}^2(r, \sigma) \end{bmatrix}, \quad (24)$$

here $q_j^2(\sigma) = a_{rj}^2 \sigma^2 + \chi_j^2$; $j = \overline{1, n+1}$.

Let's represent the integral operator H_{sn} which operates due to the formula (20) as an operator matrix-row:

$$H_{sn} [\dots] = \begin{bmatrix} \int_{R_0}^{R_1} \dots V_1(r, \lambda_s) \sigma_1 r dr & \int_{R_1}^{R_2} \dots V_2(r, \lambda_s) \sigma_2 r dr \\ \int_{R_{n-1}}^{R_n} \dots V_n(r, \lambda_s) \sigma_n r dr & \int_{R_n}^R \dots V_{n+1}(r, \lambda_s) \sigma_{n+1} r dr \end{bmatrix}. \quad (25)$$

Let's apply the operator matrix-row (25) to the problem (23), (24) according to the matrix multiplication rule. As a result of the identity (22), we get a Cauchy problem

$$\sum_{j=1}^{n+1} \left(\frac{d^2}{dt^2} + \lambda_s^2 + \gamma_j^2 + q_j^2(\sigma) \right) \tilde{u}_{jm}(t, \lambda_s, \sigma) = \sum_{j=1}^{n+1} \tilde{F}_{jm}(t, \lambda_s, \sigma) -$$

$$- a_1^2 R_0 \sigma_1 \left(\alpha_{11}^0 \right)^{-1} V_1(R_0, \lambda_s) \tilde{g}_{0m}(t, \sigma) + a_{n+1} R \sigma_{n+1} \left(\alpha_{22}^{n+1} \right)^{-1} V_{n+1}(R, \lambda_s) \tilde{g}_m(t, \sigma), \quad (26)$$

$$\sum_{j=1}^{n+1} \tilde{u}_{jm}(t, \lambda_s, \sigma) \Big|_{t=0} = \sum_{j=1}^{n+1} \tilde{g}_{jm}^1(\lambda_s, \sigma); \quad \frac{d}{dt} \sum_{j=1}^{n+1} \tilde{u}_{jm}(t, \lambda_s, \sigma) \Big|_{t=0} = \sum_{j=1}^{n+1} \tilde{g}_{jm}^2(\lambda_s, \sigma), \quad (27)$$

$$\text{here } \tilde{u}_{jm}(t, \lambda_s, \sigma) = \int_{R_{j-1}}^{R_j} \tilde{u}_{jm}(t, r, \sigma) V_j(r, \lambda_s) \sigma_j r dr; \quad j = \overline{1, n+1},$$

$$\tilde{F}_{jm}(t, \lambda_s, \sigma) = \int_{R_{j-1}}^{R_j} \tilde{F}_{jm}(t, r, \sigma) V_j(r, \lambda_s) \sigma_j r dr, \quad j = \overline{1, n+1},$$

$$\tilde{g}_{jm}^k(\lambda_s, \sigma) = \int_{R_{j-1}}^{R_j} \tilde{g}_{jm}^k(r, \sigma) V_j(r, \lambda_s) \sigma_j r dr; \quad j = \overline{1, n+1}; \quad k = 1, 2.$$

Let's suppose that $\max\{q_1^2, q_2^2, \dots, q_{n+1}^2\} = q_1^2$ and put everywhere $\gamma_j^2 = q_1^2 - q_j^2; j = \overline{1, n+1}$. Cauchy problem (26), (27) takes the form

$$\frac{d^2 \tilde{u}_m}{dt^2} + \Delta^2(\lambda_s, \sigma) \tilde{u}_m = \tilde{P}_m(t, \lambda_s, \sigma), \quad (28)$$

$$\tilde{u}_m(t, \lambda_s, \sigma) \Big|_{t=0} = \tilde{g}_m^1(\lambda_s, \sigma); \quad \frac{d\tilde{u}_m}{dt} \Big|_{t=0} = \tilde{g}_m^2(\lambda_s, \sigma), \quad (29)$$

$$\text{here } \tilde{u}_m(t, \lambda_s, \sigma) = \sum_{j=1}^{n+1} \tilde{u}_{jm}(t, \lambda_s, \sigma); \quad \tilde{F}_m(t, \lambda_s, \sigma) = \sum_{j=1}^{n+1} \tilde{F}_{jm}(t, \lambda_s, \sigma);$$

$$\tilde{g}_m^k(\lambda_s, \sigma) = \sum_{j=1}^{n+1} \tilde{g}_{jm}^k(\lambda_s, \sigma); \quad k = 1, 2; \quad \Delta^2(\lambda_s, \sigma) = \lambda_s^2 + a_{z1}^2 \sigma^2 + \chi_1^2;$$

$$\begin{aligned} \tilde{P}_m(t, \lambda_s, \sigma) = & \tilde{F}_m(t, \lambda_s, \sigma) - a_1^2 R_0 \sigma_1 \left(\alpha_{11}^0 \right)^{-1} V_1(R_0, \lambda_s) \tilde{g}_{0m}(t, \sigma) + \\ & + a_{n+1}^2 R \sigma_{n+1} \left(\alpha_{22}^{n+1} \right)^{-1} V_{n+1}(R, \lambda_s) \tilde{g}_m(t, \sigma). \end{aligned}$$

It is known [13] that the only solution of the problem (28), (29) is a function

$$\begin{aligned} \tilde{u}_m(t, \lambda_s, \sigma) = & G(t, \lambda_s, \sigma) \tilde{g}_m^2(\lambda_s, \sigma) + \frac{d}{dt} G(t, \lambda_s, \sigma) \tilde{g}_m^1(\lambda_s, \sigma) + \\ & + \int_0^t G(t-\tau, \lambda_s, \sigma) \tilde{P}_m(\tau, \lambda_s, \sigma) d\tau, \end{aligned} \quad (30)$$

here Cauchy function $G(t, \lambda_s, \sigma) = \frac{\sin(\Delta(\lambda_s, \sigma)t)}{\Delta(\lambda_s, \sigma)}$.

The superposition of operators H_{sn} and H_{sn}^{-1} is a single operator $(H_{sn} \circ H_{sn}^{-1} = H_{sn}^{-1} \circ H_{sn} = I)$. Integral operator H_{sn}^{-1} , as inverse to H_{sn} , we represent as the operator matrix-column:

$$H_{sn}^{-1} [\dots] = \begin{bmatrix} \sum_{s=1}^{\infty} \dots \frac{V_1(r, \lambda_s)}{\|V(r, \lambda_s)\|^2} \\ \sum_{s=1}^{\infty} \dots \frac{V_2(r, \lambda_s)}{\|V(r, \lambda_s)\|^2} \\ \dots \dots \dots \\ \sum_{s=1}^{\infty} \dots \frac{V_{n+1}(r, \lambda_s)}{\|V(r, \lambda_s)\|^2} \end{bmatrix}. \quad (31)$$

Let's apply operator matrix-column (31) to the matrix-element $[\tilde{u}_m(t, \lambda_s, \sigma)]$, where the function $\tilde{u}_m(t, \lambda_s, \sigma)$ is defined by formula (30) due to matrices multiplication rule. As a result we get the only solution of one-dimensional hyperbolic initial boundary problem of conjugation (16)–(19):

$$\begin{aligned} \tilde{u}_{jm}(t, r, \sigma) = & \sum_{s=1}^{\infty} \left[G(t, \lambda_s, \sigma) \tilde{g}_m^2(\lambda_s, \sigma) + \frac{\partial}{\partial t} G(t, \lambda_s, \sigma) \tilde{g}_m^1(\lambda_s, \sigma) + \right. \\ & \left. + \int_0^t G(t-\tau, \lambda_s, \sigma) \tilde{p}_m(\tau, \lambda_s, \sigma) d\tau \right] \frac{V_j(r, \lambda_s)}{\|V(r, \lambda_s)\|^2}; \quad j = \overline{1, n+1}. \end{aligned} \quad (32)$$

If to apply consistently inverse operators F_+^{-1} and F_m^{-1} to functions $\tilde{u}_{jm}(t, r, \sigma)$, which are defined by formulas (32) and perform the some simple transformation, we get functions

$$\begin{aligned} u_j(t, r, \varphi, z) = & \sum_{k=1}^{n+1} \int_0^t \int_{R_{k-1}}^{R_k} \int_0^{2\pi} \int_0^{+\infty} E_{jk}(t-\tau, r, \rho, \varphi-\alpha, z, \xi) f_k(\tau, \rho, \alpha, \xi) \times \\ & \times \sigma_k \rho d\xi d\alpha d\rho d\tau + \frac{\partial}{\partial t} \sum_{k=1}^{n+1} \int_{R_{k-1}}^{R_k} \int_0^{2\pi} \int_0^{+\infty} E_{jk}(t, r, \rho, \varphi-\alpha, z, \xi) g_k^1(\rho, \alpha, \xi) \times \\ & \times \sigma_k \rho d\xi d\alpha d\rho + \sum_{k=1}^{n+1} \int_{R_{k-1}}^{R_k} \int_0^{2\pi} \int_0^{+\infty} E_{jk}(t, r, \rho, \varphi-\alpha, z, \xi) g_k^2(\rho, \alpha, \xi) \times \\ & \times \sigma_k \rho d\xi d\alpha d\rho + \sum_{k=1}^{n+1} a_{zk}^2 \int_0^t \int_{R_{k-1}}^{R_k} \int_0^{2\pi} W_{jk}(t-\tau, r, \rho, \varphi-\alpha, z) g_k(\tau, \rho, \alpha) \times \end{aligned} \quad (33)$$

$$\begin{aligned} & \times \sigma_k \rho d\alpha d\rho d\tau + \int_0^t \int_0^{2\pi} \int_0^{+\infty} [W_{jr}^1(t-\tau, r, \varphi-\alpha, z, \xi) g_0(\tau, \alpha, \xi) + \\ & + W_{jr}^2(t-\tau, r, \varphi-\alpha, z, \xi) g(\tau, \alpha, \xi)] d\xi d\alpha d\tau. \end{aligned}$$

Functions (33) define the only solution of hyperbolic initial boundary problem of conjugation (1)–(5).

In formulas (33) there are components

$$\begin{aligned} E_{jk}(t, r, \rho, \varphi, z, \xi) &= \frac{1}{2\pi} \sum_{m=0}^{\infty} \varepsilon_m \sum_{s=1}^{\infty} \int_0^{+\infty} G(t, \lambda_s, \sigma) K(z, \sigma) K(\xi, \sigma) d\sigma \times \\ & \times \frac{V_j(r, \lambda_s) W_k(\rho, \lambda_s)}{\|V(r, \lambda_s)\|^2} \cos(m\varphi); j, k = \overline{1, n+1} \end{aligned}$$

of matrix of influence (function of influence), components $W_{jk}(t, r, \rho, \varphi, z) = E_{jk}(t, r, \rho, \varphi, z, 0)$ of tangential Green's matrix (tangential Green's function), components $W_{jr}^1(t, r, \varphi, z, \xi) = -a_1^2 R_0 \sigma_1 \left(\alpha_{11}^0\right)^{-1} E_{j1}(t, r, R_0, \varphi, z, \xi)$ of left radial Green's matrix (left radial Green's function) and components $W_{jr}^2(t, r, \varphi, z, \xi) = a_{n+1}^2 R \sigma_{n+1} \left(\alpha_{22}^{n+1}\right)^{-1} E_{j, n+1}(t, r, R, \varphi, z, \xi)$ of right radial Green's matrix (right radial Green's function) of considered problem.

Using a properties of functions of influence $E_{jk}(t, r, \rho, \varphi, z, \xi)$ and Green's functions $W_{jk}(t, r, \rho, \varphi, z)$, $W_{jr}^s(t, r, \varphi, z, \xi)$, $s = 1, 2$, we can verify that functions $u_j(t, r, \varphi, z)$ which are defined by formulas (33) satisfy the equation (1), the initial conditions (2), the boundary conditions (3), (4) and conjugate conditions (5) in the sense of theory of generalized functions [18].

The uniqueness of the solution (33) follows from its structure (integral image) and from uniqueness of the main solutions (functions of influence and Green's functions) of problem (1)–(5).

By methods from [18, 19] can be proved that under appropriate conditions on the initial data, formulas (33) define a limited classical solution of the hyperbolic initial boundary problem of conjugation (1)–(5).

We get the following theorem as the summary of the above results.

Theorem. If functions $f_j(t, r, \varphi, z)$, $g_j^1(r, \varphi, z)$, $g_j^2(r, \varphi, z)$, $g_j(t, r, \varphi)$ satisfy conditions:

- 1) are continuously differentiated twice for each variable;
- 2) have a limited variation for the geometric variables;

- 3) are absolutely summable with the variable z in $(0; +\infty)$;
- 4) conjugate conditions are true and functions $g_0(t, \varphi, z)$, $g(t, \varphi, z)$ are continuously differentiated twice for each variable, have a limited variation for the geometric variables, are absolutely summable with the variable z in $(0; +\infty)$, then hyperbolic initial boundary value problem (1)–(5) has the only limited classical solution, which is determined by formula (33).

Remark 1. In the case of $a_{rj} = a_{\varphi j} = a_{zj} \equiv a_j > 0$ formulas (33) define the structure of the solution of hyperbolic initial boundary value problem (1)–(5) in an isotropic semibounded piecewise homogeneous hollow cylinder.

Remark 2. Parameters $\alpha_{11}^0, \beta_{11}^0; \alpha_{22}^{n+1}, \beta_{22}^{n+1}$ allow to allocate the solutions of initial boundary value problems from formulas (33) in the case of boundary condition of the 1st kind, 2nd kind and 3rd kind on the radial surfaces $r = R_0$, $r = R$.

Remark 3. Parameter h allow to allocate the solutions of initial boundary value problems from formulas (33) in the case of boundary condition of the 1st kind ($h \rightarrow +\infty$) and 2nd kind ($h \rightarrow +0$) on the surface $z = 0$.

Remark 4. Analysis of the solution (33) is done directly from the general structure according to the analytical expression of functions $f_j(t, r, \varphi, z)$, $g_j^1(r, \varphi, z)$, $g_j^2(r, \varphi, z)$, $g_j(t, r, \varphi)$, $g_0(t, \varphi, z)$, $g(t, \varphi, z)$.

Remark 5. In the case of $\chi_j \equiv 0$ equation (1) is a classic three-dimensional inhomogeneous wave equation (the equation of fluctuations) for an orthotropic environment in cylindrical coordinates.

Remark 6. In the case of $\alpha_{11}^k = 0$, $\beta_{11}^k = 1$; $\alpha_{12}^k = 0$, $\beta_{12}^k = 1$; $\alpha_{21}^k = E_1^k$, $\beta_{21}^k = 0$; $\alpha_{22}^k = E_2^k$, here — Young's modulus (E), the conjugate conditions (5) coincide with conditions of ideal mechanical contact.

Thus, in these cases 5, 6 considered hyperbolic boundary value problem of mathematical physics (1)–(5) is a mathematical model of forced oscillating processes in semibounded piecewise homogeneous hollow cylinder.

Conclusions. By means of method of integral and hybrid integral transforms with the method of principal solutions (influence functions and Green's functions) integral image of exact analytical solution of hyperbolic boundary-value problem of mathematical physics in semibounded piecewise homogeneous hollow cylinder is obtained. The obtained solution is of algorithmic character, continuously depend on the parameters and data of problem and can be used in further theoretical research and in practical engineering calculations of real processes which are modeled by hyperbolic boundary-value problems of mathematical physics in piecewise homogeneous domains.

References:

1. Hadamard J. The Cauchy problem for linear partial differential equations of parabolic type / J. Hadamard. — Moscow : Nauka, 1978. — 352 p.
2. Gording L. Cauchy's problem for hyperbolic equations / L. Gording. — Moscow : IL, 1961. — 122 p.
3. Mytropol'skiy Yu. Asymptotic methods of investigation of quasi-wave equations of hyperbolic type / Yu. Mytropol'skiy, G. Khoma, M. Gromiak. — Kyiv : Naukova Dumka, 1991. — 232 p.
4. Samoilenko A. Numerical-analytic methods in the theory of periodic solutions of partial differential equations / A. Samoilenko, B. Tkach. — Kyiv : Naukova Dumka, 1992. — 208 p.
5. Smirnov M. Degenerating elliptic and hyperbolic equations / M. Smirnov. — Moscow : Nauka, 1962. — 292 p.
6. Cherniatyn V. Fourier method in mixed problem for partial differential equations / V. Cherniatyn. — Moscow : Izd. MGU, 1991. — 112 p.
7. Sergienko I. Mathematic modeling and the study of processes in heterogeneous environments / I. Sergienko, V. Skopetsky, V. Deineka. — Kyiv : Naukova Dumka, 1991. — 432 p.
8. Deineka V. Models and methods of solving of problems with conjugate conditions / V. Deineka, I. Sergienko, V. Skopetsky. — Kyiv : Naukova Dumka, 1998. — 614 p.
9. Konet I. The temperature fields in the piece-homogeneous cylindrical domains / I. Konet, M. Leniuk. — Chernivtsi : Prut, 2004. — 276 p.
10. Gromyk A. The temperature fields in the piece-homogeneous spatial environments / A. Gromyk, I. Konet, M. Leniuk. — Kamenets-Podilsky : Abetka-Svit, 2011. — 200 p.
11. Konet I. Hyperbolic boundary-value problems of mathematical physics in piecewise homogeneous spacial environments / I. Konet. — Kamenets-Podilsky : Abetka-Svit, 2013. — 120 p.
12. Konet I. Parabolic boundary value problems in piecewise homogeneous environments / I. Konet, T. Pylypiuk. — Kamenets-Podilsky : Abetka-Svit, 2016. — 244 p.
13. Gromyk A. Hyperbolic boundary value problem for semibounded piecewise-homogeneous solid cylinder / A. Gromyk, I. Konet, T. Pylypiuk // Mathematical and computer modelling. Series: Physics and Mathematics: scientific journal / Glushkov Institute of Cybernetics of National Academy of Sciences of Ukraine, Kamianets-Podilsky National Ivan Ohienko University. — Kamianets-Podilsky : Kamianets-Podil. Nat. I. Ohienko Un., 2017. — Issue 16. — P. 48–59.
14. Perestiuk M. The theory of equations of mathematical physics / M. Perestiuk, V. Marynets'. — Kyiv : Lybid', 2006. — 424 p.
15. Sneddon I. Fourier transforms / I. Sneddon. — Moscow : IL, 1955. — 668 p.
16. Tranter K. Integral transformations in mathematical physics / K. Tranter. — Moscow : Gostheorizdat, 1956. — 204 p.
17. Bybliv O. Integral Hankel transform of the 2nd kind for piecewise-homogeneous segments / O. Bybliv, M. Lenyuk // News of univ. Mathematics. — 1987. — № 5. — P. 82–85.

18. Shilov G. Mathematical analysis. Second special course / G. Shilov. — Moscow : Nauka, 1965. — 328 p.
19. Gelfand I. Some questions in the theory of differential equations / I. Gelfand, G. Shilov. — Moscow : Fizmatgiz, 1958. — 274 p.

ГІПЕРБОЛІЧНА КРАЙОВА ЗАДАЧА ДЛЯ НАПІВОБМЕЖЕНОГО КУСКОВО-ОДНОРІДНОГО ПОРОЖНИСТОГО ЦИЛІНДРА

Методом інтегральних і гібридних інтегральних перетворень у поєднанні з методом головних розв'язків (матриць впливу та матриць Гріна) вперше побудовано інтегральне зображення точного аналітичного розв'язку гіперболічної крайової задачі математичної фізики для напівобмеженого кусково-однорідного порожнистого циліндра.

Ключові слова: гіперболічне рівняння, початкові та крайові умови, умови спряження, інтегральні перетворення, головні розв'язки.

Отримано: 18.05.2018

УДК 517.5

У. В. Гудима, канд. фіз.-мат. наук,

В. О. Гнатюк, канд. фіз.-мат. наук

Кам'янець-Подільський національний університет
імені Івана Огієнка, м. Кам'янець-Подільський

КРИТЕРІЇ УЗАГАЛЬНЕНОГО ЧЕБИШОВСЬКОГО У РОЗУМІННІ ЗВАЖЕНИХ ВІДСТАНЕЙ ЦЕНТРА КІЛЬКОХ ТОЧОК ЛІНІЙНОГО НОРМОВАНОГО ПРОСТОРУ ВІДНОСНО ОПУКЛОЇ МНОЖИНИ ЦЬОГО ПРОСТОРУ

Загальновідомо, що визначальною ідеєю в питаннях зв'язків математики з практикою є ідея наближення.

Однією з центральних галузей теорії наближення є теорія наближення функцій, родоначальником якої вважається П. Л. Чебишов. У 50-х роках XIX століття він ввів поняття найкращого наближення неперервної на відріжку функції за допомогою алгебраїчних поліномів заданого степеня. Згодом було досліджено велику кількість подібних задач.

З розвитком теорії лінійних нормованих просторів стало зрозумілим, що низка задач найкращого наближення є частинними випадками задачі найкращого наближення елемента лінійного нормованого простору опуклою множиною цього простору.

Важливим питанням дослідження цієї задачі є встановлення критеріїв її екстремального елемента.

Загальний критерій екстремального елемента задачі найкращого наближення елемента лінійного нормованого простору опуклою множиною цього простору, оснований на співвід-